

SIMULTANEOUS DIOPHANTINE APPROXIMATION ON A RATIONAL ELLIPSE

LENNY FUKSHANSKY

ABSTRACT. We discuss simultaneous Diophantine approximation of points on the unit circle by rational points on the same circle, and some of its consequences. This text is included in [1].

Hlawka [2] proved the following theorem on simultaneous Diophantine approximation by quotients of Pythagorean triples.

Theorem 1. *Let $x \in (0, 1)$ be a real number. Then there exist infinitely many Pythagorean triples $(p, \sqrt{q^2 - p^2}, q)$ such that*

$$(1) \quad \left| x - \frac{p}{q} \right| \leq \frac{2\sqrt{2}}{q}.$$

Remark 1. In fact, the constant in the upper bound of Theorem 1 can be improved at least when $x = \frac{\sqrt{3}}{2}$. In [1] (Lemma 5.1) I proved that there exist infinitely many Pythagorean triples $(p, \sqrt{q^2 - p^2}, q)$ such that

$$(2) \quad \frac{1}{(2 + \sqrt{3})q} - \frac{2}{(2 + \sqrt{3})q^2} < \left| \frac{\sqrt{3}}{2} - \frac{p}{q} \right| < \frac{1}{2\sqrt{3}q},$$

explicitly constructing such an infinite sequence of Pythagorean triples.

We can use Theorem 1 to approximate points on a unit circle with rational points on the same circle.

Corollary 2. *Let (x, y) be a point on the unit circle. Then either $x, y \in \{0, \pm 1\}$, or there exist infinitely many rational points $(p/q, r/q)$ on the same circle such that*

$$(3) \quad \max \left\{ \left| x - \frac{p}{q} \right|, \left| y - \frac{r}{q} \right| \right\} \leq \frac{2\sqrt{2}}{q}.$$

Proof. First notice that it suffices to prove the statement of this corollary for the case $0 < x, y < 1$, namely the case when the point in question lies in the first quadrant, since any other point on the circle can be obtained from those in the first quadrant by a rational rotation. Let c be an arbitrary real number in the interval $(0, 1)$, then either

$$(4) \quad 0 < x \leq \sqrt{1 - c^2} < 1, \quad c \leq y < 1,$$

or

$$(5) \quad 0 < y \leq \sqrt{1 - c^2} < 1, \quad c \leq x < 1.$$

1991 *Mathematics Subject Classification.* Primary: 11J13.

Key words and phrases. simultaneous Diophantine approximation.

First assume that (4) holds. By Theorem 1, there exist infinitely many Pythagorean triples (p, r, q) with $r = \sqrt{q^2 - p^2}$ which satisfy (1). Then:

$$\begin{aligned}
 \frac{2\sqrt{2}}{q} \geq \left| x - \frac{p}{q} \right| &= \left| \sqrt{1-y^2} - \sqrt{1 - \frac{r^2}{q^2}} \right| = \frac{\left| \frac{r^2}{q^2} - y^2 \right|}{\sqrt{1-y^2} + \sqrt{1 - \frac{r^2}{q^2}}} \\
 (6) \qquad &= \frac{\frac{r}{q} + y}{\sqrt{1-y^2} + \sqrt{1 - \frac{r^2}{q^2}}} \left| y - \frac{r}{q} \right| \geq \frac{c \left(1 + \frac{n}{n+1} \right)}{2\sqrt{1 - \frac{n^2}{(n+1)^2} c^2}} \left| y - \frac{r}{q} \right|.
 \end{aligned}$$

The last inequality is true because $\frac{w+z}{\sqrt{1-w^2} + \sqrt{1-z^2}}$ is an increasing function in both variables for $0 < z, w < 1$; since $y \geq c$, we can pick q large enough so that r/q would have to be sufficiently close to y so that $r/q \geq \frac{n}{n+1}c$ for some $n \in \mathbb{Z}_{>0}$, then $r/q + y \geq c \left(1 + \frac{n}{n+1} \right)$, and $\sqrt{1-y^2} + \sqrt{1 - \frac{r^2}{q^2}} \leq 2\sqrt{1 - \frac{n^2}{(n+1)^2} c^2}$. Then (6) implies:

$$(7) \qquad \left| y - \frac{r}{q} \right| \leq \frac{\sqrt{1 - \frac{n^2}{(n+1)^2} c^2}}{c \left(1 + \frac{n}{n+1} \right)} \times \frac{4\sqrt{2}}{q}.$$

Since our choice of $c \in (0, 1)$ and positive integer n was arbitrary, we can for instance choose

$$(8) \qquad c = \frac{2n+2}{\sqrt{8n^2+4n+1}},$$

and take $n = 2$, in which case, combining (1), (7), and (8), we obtain (3).

If, on the other hand, (5) holds instead of (4), simply repeat the above argument interchanging x with y and p/q with r/q . This completes the proof. \square

A related result has also been obtained by Kopetzky in [3] (also see [4]), however his bounds are different in flavor in the sense that the constants in the upper bounds depend on x and y . Notice that the bound of Corollary 2 can be easily extended to any rational ellipse.

Corollary 3. *Let (x, y) be a point on the ellipse E , given by the equation*

$$\left(\frac{x}{a} \right)^2 + \left(\frac{y}{b} \right)^2 = 1,$$

where a, b are positive rational numbers. Then either $(x, y) = (\pm a, 0), (0, \pm b)$, or there exist infinitely many rational points $(p/q, r/q)$ on the same ellipse such that

$$(9) \qquad \max \left\{ \left| x - \frac{p}{q} \right|, \left| y - \frac{r}{q} \right| \right\} \leq \frac{2\sqrt{2} \max\{a, b\}}{q}.$$

Proof. Notice that the map $(x, y) \mapsto (x/a, y/b)$ is a bijection between E and the unit circle, which takes rational points to rational points. Now apply Corollary 2 to points of the form $(x/a, y/b)$. \square

REFERENCES

- [1] L. Fukshansky. On similarity classes of well-rounded sublattices of Z^2 . *J. Number Theory*, 129(10):2530–2556, 2009.
- [2] E. Hlawka. Approximation von irrationalzahlen und pythagoraische tripel. In *Lectures from the Colloquium on the Occasion of Ernst Peschl's 70th birthday*, volume 121, pages 1–32. Bonner Math. Schriften, Univ. Bonn, Bonn, 1980.
- [3] H. G. Kopetzky. Rationale approximationen am einheitskreis. *Monatsh. Math.*, 89(4):293–300, 1980.
- [4] H. G. Kopetzky. Diophantische approximationen auf kreisen und zyklische minima von quadratischen formen. Technical Report 179, Forschungszentrum Graz, Mathematisch-Statistische Sektion, Graz, 1981.

DEPARTMENT OF MATHEMATICS, 850 COLUMBIA AVENUE, CLAREMONT MCKENNA COLLEGE,
CA 91711

E-mail address: lenny@cmc.edu