

Heights and effective theory of quadratic forms over global fields

Lenny Fukshansky
Claremont McKenna College

Workshop on the Arithmetic of Quadratic
Forms and Integral Lattices

December 13-14, 2010

Cassels' Theorem

Let $N \geq 2$, and let

$$F(\mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \sum_{j=1}^N f_{ij} X_i Y_j$$

be a symmetric bilinear form with coefficients in \mathbb{Z} . Define its associated quadratic form

$$F(\mathbf{X}) = F(\mathbf{X}, \mathbf{X}).$$

In **1955**, **J. W. S. Cassels** proved that if F is isotropic, then there exists $\mathbf{x} \in \mathbb{Z} \setminus \{\mathbf{0}\}$ such that $F(\mathbf{x}) = 0$, and

$$(1) \quad \max_{1 \leq i \leq N} |x_i| \leq \left(3 \sum_{i=1}^N \sum_{j=1}^N |f_{ij}| \right)^{\frac{N-1}{2}}.$$

The expressions on the left and right hand sides of (1) are examples of **heights** of a vector and of a quadratic form, respectively. Since sets of integral points with explicitly bounded height are always finite, (1) provides an explicit **search bound** for non-trivial zeros of F . The exponent $\frac{N-1}{2}$ in the upper bound is sharp, as shown by an example due to **M. Kneser**.

What about heights?

The notion of height as above can be extended to number fields and function fields.

Height functions are fundamental tools of arithmetic geometry – they measure the *arithmetic complexity* of objects in much the same way that *degree* in algebraic geometry measures *geometric complexity*.

An especially important property of heights over number fields and function fields with finite coefficient fields is –

Northcott's finiteness property: For each real constant B the set of projective points with height bounded above by B is finite.

This finiteness property makes statements on the existence of objects of bounded height *effective*.

Over global fields

Analogues of Cassels' theorem over a global field K with respect to appropriately defined height functions have been obtained:

- In 1975 by **S. Raghavan** when K is a number field
- In 1987 by **A. Prestel** when K is a rational function field
- In 1997 by **A. Pfister** when K is an algebraic function field

The exponent on height of F in the upper bounds is again $\frac{N-1}{2}$.

In addition, Raghavan produced an analogous result for zeros of **hermitian forms** over CM number fields, where the exponent in the upper bound is $\frac{2N-1}{2}$.

Additional extensions of Cassels' theorem have been obtained by

- **Birch & Davenport, 1958**
- **Chalk, 1980**
- **Davenport, 1957 & 1971**
- **Schulze-Pillot, 1983**
- **Watson, 1956**

and others. In particular, results of Chalk, Davenport (1971) and Schulze-Pillot established existence theorems for collections of linearly independent isotropic vectors of bounded height.

Notation and heights

A more general effective theory followed these results. To survey it we need some notation.

Let K be either a number field, a function field, or algebraic closure of one or the other; we write \overline{K} for the algebraic closure of K , so it may be that $K = \overline{K}$. In fact, until further notice assume that $K \neq \overline{K}$.

By a function field we will always mean a finite algebraic extension of the field $\mathfrak{K} = \mathfrak{K}_0(t)$ of rational functions in one variable over a field \mathfrak{K}_0 , where \mathfrak{K}_0 can be any field.

In the number field case, we write $d = [K : \mathbb{Q}]$ for the global degree of K over \mathbb{Q} ; in the function field case, the degree is $d = [K : \mathfrak{K}]$.

Let $M(K)$ be the set of places of K . For each place $v \in M(K)$, write K_v for the completion of K at v and let d_v be the local degree of K at v , which is $[K_v : \mathbb{Q}_v]$ in the number field case, and $[K_v : \mathfrak{K}_v]$ in the function field case.

For each place u of the ground field, be it \mathbb{Q} or \mathbb{R} , we have

$$(2) \quad \sum_{v \in M(K), v|u} d_v = d.$$

If K is a number field, then for each place $v \in M(K)$ we define the absolute value $|\cdot|_v$ to be the unique absolute value on K_v that extends either the usual absolute value on \mathbb{R} or \mathbb{C} if $v|\infty$, or the usual p -adic absolute value on \mathbb{Q}_p if $v|p$, where p is a prime.

If K is a function field, then all absolute values on K are non-archimedean. For each $v \in M(K)$, let \mathcal{O}_v be the valuation ring of v in K_v and \mathfrak{M}_v the unique maximal ideal in \mathcal{O}_v . We choose the unique corresponding absolute value $|\cdot|_v$ such that:

(i) if $1/t \in \mathfrak{M}_v$, then $|t|_v = e$,

(ii) if an irreducible polynomial $p(t) \in \mathfrak{M}_v$, then $|p(t)|_v = e^{-\deg(p)}$.

In both cases, for each nonzero $a \in K$ the **product formula** reads

$$(3) \quad \prod_{v \in M(K)} |a|_v^{d_v} = 1.$$

We can now define local norms on each K_v^N :

$$|\mathbf{x}|_v = \max_{1 \leq i \leq N} |x_i|_v,$$

and for all archimedean places v also define

$$\|\mathbf{x}\|_v = \left(\sum_{i=1}^N |x_i|_v^2 \right)^{1/2},$$

for each $\mathbf{x} = (x_1, \dots, x_N) \in K_v^N$. Then define a **projective height function** on K^N by

$$H(\mathbf{x}) = \prod_{v \in M(K)} |\mathbf{x}|_v^{d_v/d}$$

for each $\mathbf{x} \in K^N$.

H is defined on the projective space $\mathbb{P}^{N-1}(K)$:

$$H(ax) = H(x), \quad \forall 0 \neq a \in K, \quad \mathbf{x} \in K^N,$$

which is true by the product formula.

For a polynomial F with coefficients in K , $H(F)$ is the height of its coefficient vector.

We also define height on subspaces of K^N . Let $V \subseteq K^N$ be an L -dimensional subspace, and let $\mathbf{x}_1, \dots, \mathbf{x}_L$ be a basis for V . Then

$$\mathbf{y} := \mathbf{x}_1 \wedge \dots \wedge \mathbf{x}_L \in K^{\binom{N}{L}}$$

under the standard embedding. Define

$$\mathcal{H}(V) := \prod_{v \nmid \infty} |\mathbf{y}|_v^{d_v/d} \times \prod_{v \mid \infty} \|\mathbf{y}\|_v^{d_v/d}.$$

This definition is independent of the choice of the basis by the product formula.

The normalizing exponent $1/d$ in the definition of H and \mathcal{H} ensures that our heights are **absolute**, i.e. do not depend on the field of definition.

Isotropic subspaces of bounded height

In **1985**, **Schlickewei** showed that if F is an isotropic rational quadratic form in $N \geq 2$ variables, and $M < N$ is the dimension of a maximal totally isotropic subspace of the quadratic space (\mathbb{Q}^N, F) , then there exists such a subspace W with

$$\mathcal{H}(W) \ll_N H(F)^{\frac{N-M}{2}}.$$

In **1987**, **Vaaler** generalized this result and extended it to number fields. Specifically, let K be a number field, $N \geq 2$ and $1 \leq L \leq N$ integers, $V \subseteq K^N$ an L -dimensional subspace, F a quadratic form over K in N variables, and (V, F) a quadratic space. Let $M \leq L/2$ be the Witt index of (V, F) and λ dimension of the radical. Then there exists a maximal totally isotropic subspace $W \subset V$ of dimension $M + \lambda$ such that

$$\mathcal{H}(W) \ll_{K,L,M,\lambda} H(F)^{\frac{L-M-\lambda}{2}} \mathcal{H}(V).$$

Analogous result over $\overline{\mathbb{Q}}$

For quadratic spaces over $\overline{\mathbb{Q}}$, we have:

(F., 2008) Let $V \subseteq \overline{\mathbb{Q}}^N$, $\dim V = L$, and consider the quadratic space (V, F) . Let V^\perp be the radical of V , so that $\dim V^\perp = \lambda$, and $M = \left\lfloor \frac{L-\lambda}{2} \right\rfloor$ is the Witt index of V . Then

$$V = V^\perp \perp U,$$

where U is regular, $\dim U = L - \lambda$, and

$$\mathcal{H}(V^\perp) \ll_L H(F)^{L-\lambda} \mathcal{H}(V)^2, \quad \mathcal{H}(U) \ll_L \mathcal{H}(V).$$

There also exists a maximal totally isotropic subspace W of U so that $\dim W = M$ and

$$\mathcal{H}(W) \ll_M \begin{cases} H(F)^{\frac{M^2}{2}} \mathcal{H}(U)^{\frac{M^2+M+2}{2M}} & \text{if } 2 \mid L - \lambda \\ H(F)^{M^2} \mathcal{H}(U)^{\frac{4M}{3}} & \text{if } 2 \nmid L - \lambda. \end{cases}$$

Then $W' := \text{span}_{\overline{\mathbb{Q}}}\{V^\perp, W\}$ is a maximal totally isotropic subspace of V , $\dim W' = M + \lambda$, and

$$\mathcal{H}(W') \leq \mathcal{H}(V^\perp) \mathcal{H}(W).$$

Families of isotropic subspaces

In **1989**, **Vaaler** proved the following result over number fields, building on previous work by **Schlickewei and Schmidt (1987)** over \mathbb{Q} :

Given a quadratic space (V, F) over a number field K , as above, there exists a family

$$U_0, \dots, U_{L-M} \subseteq V$$

of maximal totally isotropic subspaces of V such that

$$V = \text{span}_K \{W_0, \dots, W_{L-M}\}$$

and for each $0 \leq i \leq L - M$,

$$\mathcal{H}(W_0)\mathcal{H}(W_i) \ll_{K,L,M} H(F)^{L-M} \mathcal{H}(V)^2.$$

In joint work (**W. K. Chan, L. F., G. Henshaw; in progress**), we show that in fact there exists an infinite sequence of families of maximal totally isotropic subspaces

$$\{W_{n1}, \dots, W_{nJ}\}_{n=1}^{\infty} \subseteq V,$$

where $J = M(L - 2M - \lambda)$, such that for each $n \geq 1$,

$$\text{span}_K \{W_{n1}, \dots, W_{nJ}\} = V,$$

and for each $1 \leq j \leq J$,

$$\mathcal{H}(W_{nj}) \ll H(F)^{p(L,M)} \mathcal{H}(V)^{q(M)},$$

where the constant in the upper bound depends on K, N, L, M, λ, n , and $p(L, M)$, $q(M)$ are polynomials: $p(L, M)$ is linear in L , quartic in M , and $q(M)$ is cubic in M .

We have also obtained an analogous result over $\overline{\mathbb{Q}}$, and are working on the function field case.

Witt decomposition (F., 2007-2008)

Let K be a number field or $\overline{\mathbb{Q}}$, and let (V, F) be an L -dimensional quadratic space in N variables over K , as above. Let $r := L - \lambda$ be the rank of F on V . There exists an orthogonal decomposition of the quadratic space (V, F) of the form

$$V = V^\perp \perp \mathbb{H}_1 \perp \cdots \perp \mathbb{H}_M \perp U,$$

where $\mathbb{H}_1, \dots, \mathbb{H}_M$ are hyperbolic planes and U is anisotropic component, with all components of bounded height. Specifically, if K is a number field, then

$$\mathcal{H}(V^\perp) \ll_{K,L,r} H(F)^{r/2} \mathcal{H}(V),$$

as proved by **Vaaler (1989)**, and

$$\max\{\mathcal{H}(\mathbb{H}_i), \mathcal{H}(U)\} \ll \left\{ H(F)^{\frac{L+2M}{4}} \mathcal{H}(V) \right\}^{e_1(M)},$$

for each $1 \leq i \leq M$, where

$$e_1(M) = \frac{(M+1)(M+2)}{2}$$

and the constant depends on N, L, M .

If $K = \overline{\mathbb{Q}}$, then

$$\mathcal{H}(V) \ll_{L,r} H(F)^r \mathcal{H}(V)^2,$$

and

$$\mathcal{H}(\mathbb{H}_i) \ll \left\{ H(F)^{M^2+1} \mathcal{H}(V)^{\frac{6M+5}{4M+2}} \right\}^{e_2(M)},$$

where

$$e_2(M) = \frac{(M+1)(M+2)}{2} \left(\frac{3}{2} \right)^M$$

and the constant depends on L, M, r , and $U = \{0\}$ if $L = 2M$, or $U = \overline{\mathbb{Q}}\mathbf{y}$ with

$$\mathcal{H}(U) \ll_{L,M,r} \mathcal{H}(V)^{\frac{2M+3}{4M+2}},$$

if $L = 2M + 1$.

In a joint project (**W. K. Chan, F., G. Henshaw; in progress**) we are working to obtain function field analogue of these results.

Symplectic spaces (F., 2009)

Let K be a number field, function field, or the algebraic closure of one or the other. Let

$$F(\mathbf{X}, \mathbf{Y}) = \sum_{i=1}^N \sum_{j=1}^N f_{ij} X_i Y_j$$

be an alternating bilinear form in $N \geq 2$ variables with coefficients in K . Let $V \subseteq K^N$ be a $2M$ -dimensional subspace, $1 \leq M \leq N/2$, such that (V, F) is a regular symplectic space over K . Define

$$a(M) = \begin{cases} \frac{M^2+4M}{4} & \text{if } 2 \mid M \\ \frac{M^2+4M-1}{4} & \text{if } 2 \nmid M, \end{cases}$$

and

$$b(M) = \begin{cases} \frac{2M^3+9M^2-14M}{12} & \text{if } 2 \mid M \\ \frac{2M^3+9M^2-14M+3}{12} & \text{if } 2 \nmid M. \end{cases}$$

For each $1 \leq n \leq M$, there exist totally isotropic subspaces U_n and W_n of (V, F) such that

$$\dim_K U_n = \dim_K W_n = n, \quad U_n \cap W_n = \{\mathbf{0}\}$$

and

$$U_1 \subset U_2 \subset \cdots \subset U_M, \quad W_1 \subset W_2 \subset \cdots \subset W_M,$$

so that $V = \text{span}_K \{U_M, W_M\}$, and

$$\mathcal{H}(U_n)\mathcal{H}(W_n) \ll_{K,N,M} \left(\mathcal{H}(V)^{a(M)} \mathcal{H}(F)^{b(M)} \right)^{\frac{n}{M}}.$$

Moreover,

$$V = \mathbb{H}_1 \perp \cdots \perp \mathbb{H}_M,$$

where for each $1 \leq i \leq M$, \mathbb{H}_i is a hyperbolic plane, and

$$\prod_{i=1}^M \mathcal{H}(\mathbb{H}_i) \ll_{K,N,M} \mathcal{H}(V)^{a(M)} \mathcal{H}(F)^{b(M)}.$$

Hermitian forms (Chan, F., 2010)

Let K be a **totally real** number field of degree d , and let $\alpha, \beta \in O_K$ be **totally negative**. Let $D = \left(\begin{smallmatrix} \alpha, \beta \\ K \end{smallmatrix} \right)$ be a positive definite quaternion algebra over K , generated by the elements i, j, k which satisfy the relations $i^2 = \alpha$, $j^2 = \beta$, $ij = -ji = k$, $k^2 = -\alpha\beta$. Let \mathcal{O} be an order in D . Let $N \geq 2$ be an integer, and let $V \subseteq D^N$ be an L -dimensional right D -subspace, $1 \leq L \leq N$. Let $F(\mathbf{X}, \mathbf{Y}) \in D[\mathbf{X}, \mathbf{Y}]$ be a hermitian form in $2N$ variables, and assume that F is isotropic on V . Then there exists a basis $\mathbf{y}_1, \dots, \mathbf{y}_L$ for V over D such that $F(\mathbf{y}_n) := F(\mathbf{y}_n, \mathbf{y}_n) = 0$ for all $1 \leq n \leq L$, $h(\mathbf{y}_1) \leq \dots \leq h(\mathbf{y}_L)$, and

$$h(\mathbf{y}_1)h(\mathbf{y}_L) \ll_{K, \mathcal{O}, N, L, \alpha, \beta} H_{\text{inf}}(F)^{4L-1} H^{\mathcal{O}}(V)^8,$$

for appropriately defined height functions h , H_{inf} , and $H^{\mathcal{O}}$, following the approach of **C. Liebendorfer (2004)**.

This is an analogue of a result of **Vaaler (1989)** over number fields.

Further work

Some of the additional work done includes investigation of small zeros of quadratic forms over a field K outside of unions of subspaces and varieties:

- In 1998 by **D. Masser**: $K = \mathbb{Q}$, outside of one hyperplane
- In 2004 by **L. F.**: K is a number field, union of m hyperplanes
- In 2009 by **R. Dietmann**: better bound when $m > 1$
- In progress by **W. K. Chan, L. F., G. Henshaw**: K is a number field or $\overline{\mathbb{Q}}$, in a vector space missing a union of varieties

In a joint project (**W. K. Chan, F., G. Henshaw; in progress**) we are working to obtain function field analogue of these results.